

Families of exact solutions to Vasiliev's 4D equations with spherical, cylindrical and biaxial symmetry

Carlo IAZEOLLA

Università di Bologna and INFN, Sezione di Bologna

*C.I. , P. Sundell – JHEP 12 (2011) 084
arXiv:1107.1217 [hep-th]*

Summary

- The 4D Vasiliev equations
 - Oscillator algebras
 - Full equations (bosonic)
- Solving the equations
 - Gauge function method and separation of variables in twistor space
- Exact solutions
 - Projector Ansätze and six infinite families of solutions.
 - Weyl curvatures and deformed oscillators.
 - Spherically and cylindrically-symmetric solutions.
 - Construction of some HS invariants. Singularities?
- Conclusions and Outlook

Oscillator algebra

- Commuting variables $\underline{Y}_\alpha = (y_\alpha, \bar{y}_{\dot{\alpha}})$, $\underline{Z}_\alpha = (z_\alpha, -\bar{z}_{\dot{\alpha}}) \rightarrow \mathfrak{sp}(4, \mathbb{R})$ quartets

$$[Y_\alpha, Y_\beta]_\star = 2iC_{\alpha\beta} = 2i \begin{pmatrix} \varepsilon_{\alpha\beta} & 0 \\ 0 & \varepsilon_{\dot{\alpha}\dot{\beta}} \end{pmatrix}, \quad [Z_\alpha, Z_\beta]_\star = -2iC_{\alpha\beta}, \quad [Y_\alpha, Z_\beta]_\star = 0$$

- Star-product, normal-ordering (wrt $A^+ = (Y-Z)/2i$, $A^- = (Y+Z)/2$)

$$\widehat{F}(Y, Z) \star \widehat{G}(Y, Z) = \int_{\mathcal{R}} \frac{d^4U d^4V}{(2\pi)^4} e^{iV^\alpha U_\alpha} \widehat{F}(Y + U, Z + U) \widehat{G}(Y + V, Z - V)$$

- π automorphism generated by the inner kleinian operator κ :

$$\pi(\widehat{f}(y, \bar{y}; z, \bar{z})) = \widehat{f}(-y, \bar{y}; -z, \bar{z}), \quad \bar{\pi}(\widehat{f}(y, \bar{y}; z, \bar{z})) = \widehat{f}(y, -\bar{y}; z, -\bar{z})$$

$$\begin{aligned} \pi(\widehat{f}) &= \kappa \star \widehat{f} \star \kappa, & \kappa &= e^{iy^\alpha z_\alpha}, & \kappa \star \kappa &= 1 \\ \kappa &= \kappa_y \star \kappa_z, & \kappa_y \star \kappa_y &= 1 \text{ idem } \kappa_z, \bar{\kappa}_{\bar{y}} \text{ and } \bar{\kappa}_{\bar{z}} \\ \kappa_y &= 2\pi\delta^2(y) = 2\pi\delta(y_1)\delta(y_2) \end{aligned}$$

- Fields live on correspondence space, locally $\mathcal{X} \times \mathcal{Y} \times \mathcal{Z}$:

$$d \rightarrow \widehat{d} = d + d_Z = dx^\mu \frac{\partial}{\partial x^\mu} + dz^\alpha \frac{\partial}{\partial z^\alpha} + d\bar{z}^{\dot{\alpha}} \frac{\partial}{\partial \bar{z}^{\dot{\alpha}}}$$

$$A(x|Y) \rightarrow \widehat{A}(x|Z, Y) \equiv (dx^\mu \widehat{A}_\mu + dz^\alpha \widehat{A}_\alpha + d\bar{z}^{\dot{\alpha}} \widehat{A}_{\dot{\alpha}})(x|Z, Y), \quad A_\mu(x|Y) = \widehat{A}_\mu|_{Z=0}$$

$$\Phi(x|Y) \rightarrow \widehat{\Phi}(x|Z, Y), \quad \Phi(x|Y) = \widehat{\Phi}(x|Z, Y)|_{Z=0}$$

The Vasiliev Equations

- Gauge field $\in \text{Adj}(\mathfrak{hs}(3,2))$ (*master 1-form connection*):

$$A_\mu(x|y, \bar{y}) = \sum_{n+m=2\text{mod}4}^{\infty} \frac{i}{2n!m!} dx^\mu A_\mu^{\alpha_1 \dots \alpha_n \dot{\alpha}_1 \dots \dot{\alpha}_m}(x) y_{\alpha_1} \dots y_{\alpha_n} \bar{y}_{\dot{\alpha}_1} \dots \bar{y}_{\dot{\alpha}_m}$$

(every spin- s sector contains all one-form connections that are necessary for a frame-like formulation of HS dynamics (finitely many))

Generators of $\mathfrak{hs}(3,2)$: $T_s \sim y_{\alpha_1} \dots y_{\alpha_n} \bar{y}_{\dot{\alpha}_1} \dots \bar{y}_{\dot{\alpha}_m}$, $\frac{n+m}{2} + 1 = s$

Bilinears in osc. $\rightarrow \mathfrak{so}(3,2)$: $M_{AB} = -\frac{1}{8} Y^\alpha (\Gamma_{AB})_{\underline{\alpha}\beta} Y^\beta = \{M_{ab}, P_a\}$

- **Massless UIRs** with all spins in AdS include **a scalar!**

\rightarrow “twisted adjoint” *master 0-form* (contains scalar, Weyl, HS Weyl and derivatives)

$$T(X)(\Phi) = [X, \Phi]_{\star, \pi} \equiv X \star \Phi - \Phi \star \pi(X)$$

- *Weyl 0-form* : $\Phi(x|y, \bar{y}) = \sum_{|n-m|=0\text{mod}4}^{\infty} \frac{1}{n!m!} \Phi^{\alpha_1 \dots \alpha_n \dot{\alpha}_1 \dots \dot{\alpha}_m}(x) y_{\alpha_1} \dots y_{\alpha_n} \bar{y}_{\dot{\alpha}_1} \dots \bar{y}_{\dot{\alpha}_m}$

N.B.: spin- s sector \rightarrow infinite-dimensional

(upon constraints, all on-shell-nontrivial covariant derivatives of the physical fields₄ *i.e.*, all the local dof encoded in the 0-form at a point)

The Vasiliev Equations

- Full eqs: $\hat{F} \equiv \hat{d}\hat{A} + \hat{A} \star \hat{A} = \frac{i}{4} (dz^\alpha \wedge dz_\alpha \hat{\mathcal{B}} \star \hat{\Phi} \star \kappa + d\bar{z}^{\dot{\alpha}} \wedge d\bar{z}_{\dot{\alpha}} \hat{\bar{\mathcal{B}}} \star \hat{\Phi} \star \bar{\kappa})$
(Vasiliev '90)
 $\hat{\mathcal{D}}\hat{\Phi} \equiv \hat{d}\hat{\Phi} + \hat{A} \star \hat{\Phi} - \hat{\Phi} \star \bar{\pi}(\hat{A}) = 0$

Local sym: $\delta\hat{A} = \hat{D}\hat{\epsilon} , \quad \delta\hat{\Phi} = -[\hat{\epsilon}, \hat{\Phi}]_\pi$

- In components: $\hat{F}_{\mu\nu} = \hat{F}_{\mu\alpha} = \hat{F}_{\mu\dot{\alpha}} = 0 , \quad \hat{D}_\mu\hat{\Phi} = 0 ,$
 $[\hat{S}_\alpha, \hat{S}_\beta]_\star = -2i\epsilon_{\alpha\beta}(1 - \mathcal{B} \star \hat{\Phi} \star \kappa) ,$
 $[\hat{S}_{\dot{\alpha}}, \hat{S}_{\dot{\beta}}]_\star = -2i\epsilon_{\dot{\alpha}\dot{\beta}}(1 - \bar{\mathcal{B}} \star \hat{\Phi} \star \bar{\kappa})$
 $[\hat{S}_\alpha, \hat{S}_{\dot{\beta}}]_\star = 0 ,$
 $\hat{S}_\alpha \star \hat{\Phi} + \hat{\Phi} \star \pi(\hat{S}_\alpha) = 0 ,$
 $\hat{S}_{\dot{\alpha}} \star \hat{\Phi} + \hat{\Phi} \star \bar{\pi}(\hat{S}_{\dot{\alpha}}) = 0$
 $\hat{S}_\alpha = z_\alpha - 2i\hat{A}_\alpha$

- Z-evolution determines Z-contractions in terms of original dof.
Solution of Z-eqs. yields consistent nonlinear corrections as an expansion in Φ .

Black Holes and Higher Spins

- Crucial to look into the non-perturbative sector of the theory, may shed some light on peculiarities of HS physics and prompts to study global issues in HS gravity (boundary conditions, asymptotic charges, global dof in $\mathcal{Z}...$). Very likely new tools, and HS geometry adapted to HS symmetries, have to be developed.
- HS Gravity does not admit a consistent truncation to spin 2. No obvious embedding of gravitational bhs.
- Characterization of bhs rests on geodesic motion, but relativistic interval $ds^2 = g_{\mu\nu} dx^\mu dx^\nu$ is NOT HS-invariant . What is to be called a “higher-spin black hole”?
- Do non-local interactions & HS gauge symmetries smooth out singularities? (already from ST we are used to higher-derivative stringy correction affecting the nature of singularities)

Exact solutions: gauge function method

- Y x Z-space eqns:

$$\begin{aligned}
 \widehat{F}_{\mu\nu} &= \widehat{F}_{\mu\alpha} = \widehat{F}_{\mu\dot{\alpha}} = 0, & \widehat{D}_\mu \widehat{\Phi} &= 0, \\
 [\widehat{S}'_\alpha, \widehat{S}'_\beta]_\star &= -2i\epsilon_{\alpha\beta}(1 - \mathcal{B} \star \widehat{\Phi}' \star \kappa), \\
 [\widehat{S}'_{\dot{\alpha}}, \widehat{S}'_{\dot{\beta}}]_\star &= -2i\epsilon_{\dot{\alpha}\dot{\beta}}(1 - \bar{\mathcal{B}} \star \widehat{\Phi}' \star \bar{\kappa}) \\
 [\widehat{S}'_\alpha, \widehat{S}'_{\dot{\beta}}]_\star &= 0, \\
 \widehat{S}'_\alpha \star \widehat{\Phi}' + \widehat{\Phi}' \star \pi(\widehat{S}'_\alpha) &= 0, \\
 \widehat{S}'_{\dot{\alpha}} \star \widehat{\Phi}' + \widehat{\Phi}' \star \bar{\pi}(\widehat{S}'_{\dot{\alpha}}) &= 0
 \end{aligned}$$

- Project on \mathcal{Z} ! (base \leftrightarrow fiber evolution)

Locally give x-dep. via gauge functions (spacetime \sim pure gauge!)

$$\widehat{A}_\mu = \widehat{L}^{-1} \star \partial_\mu \widehat{L}, \quad \widehat{S}_\alpha = \widehat{L}^{-1} \star (\widehat{S}'_\alpha) \star \widehat{L}, \quad \widehat{\Phi} = \widehat{L}^{-1} \star \widehat{\Phi}' \star \pi(\widehat{L})$$

$$\widehat{L} = \widehat{L}(x|Z, Y), \quad \widehat{L}(0|Z, Y) = 1 \quad \widehat{S}'_\alpha = \widehat{S}_\alpha(0|Z, Y), \quad \widehat{\Phi}' = \widehat{\Phi}(0|Z, Y)$$

- Z-eq.^{ns} can be solved exactly: 1) imposing **symmetries** on primed fields
2) via **projectors**
- “Dress” with x-dependence by performing star-products with gauge function.

AdS₄ Vacuum Solution

- AdS₄ vacuum sol.:

$$\widehat{\Phi} = 0, \quad \widehat{S}_\alpha = \widehat{S}_\alpha^{(0)} = z_\alpha, \quad \widehat{S}_{\dot{\alpha}} = \widehat{S}_{\dot{\alpha}}^{(0)} = \bar{z}_{\dot{\alpha}}, \quad \widehat{A}_\mu = \Omega_\mu^{(0)} = L^{-1} \star \partial_\mu L$$

The gauge function

$$(h = \sqrt{1 - \lambda^2 x^2})$$

$$L(x; y, \bar{y}) = e_\star^{i\lambda \tilde{x}^\mu(x) \delta_\mu^a P_a} = \frac{2h}{1+h} \exp \left[\frac{i\lambda x^{\alpha\dot{\alpha}} y_\alpha \bar{y}_{\dot{\alpha}}}{1+h} \right]$$

gives AdS₄ connection

$$\Omega_\mu^{(0)} = -i \left(\frac{1}{2} \omega_{(0)}^{ab} M_{ab} + e_{(0)}^a P_a \right) = \frac{1}{4i} \left(\omega_{(0)}^{\alpha\beta} y_\alpha y_\beta + \bar{\omega}_{(0)}^{\dot{\alpha}\dot{\beta}} \bar{y}_{\dot{\alpha}} \bar{y}_{\dot{\beta}} + 2e_{(0)}^{\alpha\dot{\beta}} y_\alpha \bar{y}_{\dot{\beta}} \right)$$

$$e_{(0)}^{\alpha\dot{\alpha}} = -\frac{\lambda(\sigma^a)^{\alpha\dot{\alpha}} dx_a}{h^2}, \quad \omega_{(0)}^{\alpha\beta} = -\frac{\lambda^2(\sigma^{ab})^{\alpha\beta} dx_a dx_b}{h^2}$$

leading to AdS₄ metric in stereographic coords.:

$$ds_{(0)}^2 = \frac{4dx^2}{(1 - \lambda^2 x^2)^2}$$

- Global symmetries:

$$\delta S_\alpha^{(0)} = [z_\alpha, \widehat{\epsilon}]_\star = 0 \Rightarrow \widehat{\epsilon} = \epsilon^{(0)}(x|Y)$$

$$\delta \Omega_\mu^{(0)} = D_\mu^{(0)} \epsilon^{(0)}(x|Y) = 0$$

Y²-sector: $\epsilon^{(0)} = -i \left(\frac{1}{2} \kappa^{ab} M_{ab} + v^a P_a \right)$



$$\delta e_{(0)}^a = 0 \Rightarrow \nabla_a^{(0)} v_b = \kappa_{ab}$$

$$\delta \omega_{(0)}^{ab} = 0 \Rightarrow \nabla_a^{(0)} \kappa_{bc} = g_{ac}^{(0)} v_b - g_{ab}^{(0)} v_c$$

Local properties of 4D black holes

- Bh Weyl tensor is of **Petrov-type D**, ((anti-)selfdual part) has 2 principal spinors :

$$\Phi_{\alpha\beta\gamma\delta} = \nu(x) u_{(\alpha}^+ u_{\beta}^- u_{\gamma}^+ u_{\delta)}^-, \quad u^{+\alpha} u_{\alpha}^- = 1$$

- Local characterization of 4D bhs: sol.n.s of Einstein's eqs. in vacuum (flat or AdS) such that their Weyl tensor's principal spinors are collinear with those of the **Killing 2-form** of an asymptotically *timelike* KVF, $\kappa_{\mu\nu} = \nabla_{\mu} v_{\nu}$ (Mars, '99;

Didenko-Matveev-Vasiliev, '08-'09),

$$\Phi_{\alpha\beta\gamma\delta} \sim \frac{M}{(\kappa^2)^{5/2}} \kappa_{(\alpha\beta} \kappa_{\gamma\delta)}, \quad \kappa^2 := \frac{1}{2} \kappa^{\alpha\beta} \kappa_{\alpha\beta}$$

- A generic bh is completely determined by a chosen **background** global symmetry parameter $Y^{\alpha} K_{\alpha\beta} Y^{\beta}$ (Didenko-Matveev-Vasiliev, '09)

$$K_{\alpha\beta} = \begin{pmatrix} \kappa_{\alpha\beta} & v_{\alpha\dot{\beta}} \\ \bar{v}_{\dot{\alpha}\beta} & \bar{\kappa}_{\dot{\alpha}\dot{\beta}} \end{pmatrix}, \quad D_0 K_{\alpha\beta} = 0$$

Properties of bh encoded in algebraic conditions: $K^2 = -1 \rightarrow$ static:

$$K_{\alpha}^{\beta} K_{\beta}^{\gamma} = -\delta_{\alpha}^{\gamma} \Leftrightarrow \begin{cases} \kappa^2 + v^2 = 1 \\ \kappa^2 = \bar{\kappa}^2 \\ \kappa_{\alpha}^{\beta} v_{\beta}^{\dot{\gamma}} + v_{\alpha}^{\dot{\beta}} \bar{\kappa}_{\dot{\beta}}^{\dot{\gamma}} = 0 \end{cases} \longrightarrow v_{[\mu} \nabla_{\nu} v_{\rho]} = 0$$

HS black-hole-like Ansatz

- Weyl zero-form $\widehat{\Phi} = \widehat{L}^{-1} \star \widehat{\Phi}' \star \pi(\widehat{L})$: reduces eqs. to linearized on AdS

$$\partial_\mu \widehat{\Phi} + [\widehat{A}_\mu, \widehat{\Phi}]_\pi = 0 \quad \rightarrow \quad \partial_\mu \Phi + [\Omega_\mu^{(0)}, \Phi]_\pi = 0 \quad \text{with}$$

$$\widehat{L}(x|Y, Z) = L(x|Y) \star \widetilde{L}(x|Z), \quad \pi(\widetilde{L}) = \widetilde{L}; \quad \widehat{\Phi}' = \Phi'(Y)$$

- Link with global sym parameters: to any HS global sym parameter $\epsilon_{(0)}(x|Y)$ ($D^{(0)}\epsilon^{(0)} = 0$) is associated a solution $\epsilon^{(0)} \star \kappa_y$ of the linearized Weyl 0-form eqn.

$$\partial_\mu(\epsilon^{(0)} \star \kappa_y) + [\Omega_\mu^{(0)}, (\epsilon^{(0)} \star \kappa_y)]_\pi = (D^{(0)}\epsilon^{(0)}) \star \kappa_y = 0$$

$$\Phi(x|Y) = \epsilon^{(0)}(x|Y) \star \kappa_y = L^{-1} \star \epsilon'_{(0)}(Y) \star L \star \kappa_y \quad \Rightarrow \quad \Phi'(Y) = \epsilon'_{(0)}(Y) \star \kappa_y$$

- Bh determined by a chosen AdS KVF $K_{\underline{\alpha}\underline{\beta}}(x) \rightarrow$ by a rigid $K'_{\underline{\alpha}\underline{\beta}} \in \mathfrak{sp}(4, \mathbb{C})$.
Generalize to a HS global sym parameter (*Didenko-Vasiliev '09*)

$$\epsilon_0(x|Y) = f(Y^\alpha K_{\underline{\alpha}\underline{\beta}}(x) Y^\beta), \quad \Rightarrow \quad \epsilon'_0(Y) = f(Y^\alpha K'_{\underline{\alpha}\underline{\beta}} Y^\beta),$$

- Assume two commuting $K^{(+)}_{\underline{\alpha}\underline{\beta}}(x)$ and $K^{(-)}_{\underline{\alpha}\underline{\beta}}(x)$, $[K^{(+)}, K^{(-)}]_{\underline{\alpha}\underline{\beta}} = 0$.

Rigid elements $K'^{(\pm)} := Y^\alpha K'_{\underline{\alpha}\underline{\beta}} Y^\beta$ generate $\mathfrak{so}(2)_{(+)} \oplus \mathfrak{so}(2)_{(-)}$.

HS black-hole-like Ansatz

- Which $f(K')$? Choose **projectors** (enforce Kerr-Schild property in gauge fields):
expand all fluctuation fields Φ', A'_α in projectors $P_{n_1 n_2}(K'_{(+)}, K'_{(-)})$

$$P_{n_1, n_2} \star P_{n'_1, n'_2} = \delta_{n_1 n'_1} \delta_{n_2 n'_2} P_{n_1, n_2}, \quad (w_i - n_i) \star P_{n_1, n_2} = 0,$$

$$K'_{(q)} := \frac{1}{2} (w_2 + q w_1), \quad \mathbf{n} = (n_1, n_2) \in (\mathbb{Z} + \frac{1}{2}) \times (\mathbb{Z} + \frac{1}{2})$$

$$P_{n_1 n_2}(K'_{(+)}, K'_{(-)}) = 4(-1)^{|n|-1} e^{-2(w_1+w_2)} L_{n_1-\frac{1}{2}}(4w_1) L_{n_2-\frac{1}{2}}(4w_2)$$

axisymmetric excitations of vacuum with enhanced sym $\mathfrak{c}_{\mathfrak{sp}(4, \mathbb{R})}(K^{(q)})$

$$P_{1/2, 1/2}(K'_{(q)}) := 4e^{-\frac{1}{2} Y^\alpha \underline{K}'_{\alpha\beta} Y^\beta}, \quad P_{1/2, 1/2} \star P_{1/2, 1/2} = P_{1/2, 1/2}, \quad \left(\underline{K}'_{\alpha}{}^\beta \underline{K}'_{\beta}{}^\gamma = -\delta_{\alpha}{}^\gamma \right)$$

$$\underline{K}'_{\alpha}{}^\beta \underline{K}'_{\beta}{}^\gamma = -\delta_{\alpha}{}^\gamma \Rightarrow \underline{K}'_{\alpha\beta} \sim (\Gamma_{AB})_{\alpha\beta}, \quad M_{AB} = -\frac{1}{8} Y^\alpha (\Gamma_{AB})_{\alpha\beta} Y^\beta$$

- 3 inequivalent embeddings of $\mathfrak{so}(2)_{(+)} \oplus \mathfrak{so}(2)_{(-)}$ in $\mathfrak{sp}(4, \mathbb{C})$:
(E, J) ; **(J, iB)** ; **(iB, iP)**. [E:=M_{0'0}=P₀; J:=M₁₂; B:=M₀₃; P:=P₁]
- Each gives rise to two families ($|K'_{(+)}| > |K'_{(-)}|$ or $|K'_{(+)}| < |K'_{(-)}|$) based on choice of “principal” $K'_{(q)}$, determining symmetries of vacuum (and behaviour of Weyl tensors) \rightarrow six infinite families of solutions.

HS black-hole-like Ansatz

- Specific combinations of $P_{n_1 n_2} (K'_{(+)}, K'_{(-)})$ give rank- $|n|$ projectors depending on $K'_{(+)}$ only \rightarrow enhanced sym under $\mathfrak{c}_{\text{sp}(4, \mathbb{R})}(K^{(q)})$

$$\begin{aligned} \mathcal{P}_n(K'_{(q)}) &= \sum_{n_2 \epsilon_1 \epsilon_2 \bar{q} \bar{n}} P_{n_1, n_2} = 4(-)^{n - \frac{1+\epsilon}{2}} e^{-4K'_{(q)}} L_{n-1}^{(1)}(8K'_{(q)}) \\ &= 2(-)^{n - \frac{1+\epsilon}{2}} \oint_{C(\epsilon)} \frac{d\eta}{2\pi i} \left(\frac{\eta+1}{\eta-1} \right)^n e^{-4\eta K'_{(q)}}, \quad n \in \mathbb{Z} \end{aligned}$$

- $\Rightarrow \Phi'(Y) =$ any $f(Y)$ diagonalizable on such bases of projectors $\star \kappa_y$:

$$\Phi'(Y) = \sum_{\mathbf{n}} \nu_{\mathbf{n}} P_{\mathbf{n}}(Y) \star \kappa_y$$

- Weyl 0-form: $\Phi(x|Y) = \sum_n \nu_n \mathcal{N}_n \oint_{C(\epsilon)} \frac{d\eta}{2\pi i} \left(\frac{\eta+1}{\eta-1} \right)^n \underbrace{L^{-1}(x) \star e^{-4\eta K'_{(q)}} \star L(x)} \star \kappa_y$

Type-D

Weyl 0-form
generating f :

$$\bar{y} = 0: \frac{1}{\eta \sqrt{\mathcal{K}^2(x)}} \exp\left(\frac{1}{2\eta} y^\alpha \mathcal{K}_{\alpha\beta}^{-1}(x) y^\beta\right), \quad \mathcal{K}_{\alpha\beta}^{-1} = -\frac{\mathcal{K}_{\alpha\beta}}{\mathcal{K}^2}, \quad \mathcal{K}^2 = \frac{1}{2} \mathcal{K}^{\alpha\beta} \mathcal{K}_{\alpha\beta}$$

$$\Phi_{\alpha(2s)}^{(n)} \sim \frac{\nu_n}{(\mathcal{K}^2)^{s+1/2}} \mathcal{K}_{(\alpha_1 \alpha_2 \dots \alpha_{2s-1} \alpha_{2s})}$$

Spherically symmetric type-D solutions

- Based on enhanced E-dependent projectors, s.t. residual symmetry \rightarrow centralizer of E $\Rightarrow \mathfrak{e} = \mathfrak{so}(2)_E \oplus \mathfrak{so}(3)_{M_{ij}}$.

$$\delta\Phi(x|Y) = -[\epsilon(x|Y), \Phi(x|Y)]_{\star, \pi} = 0 \Leftrightarrow [\epsilon'(Y), e^{-4sK_{(q)}}]_{\star} = 0 \Rightarrow K_{(q)} = E$$

- Spherical symm. solutions based on scalar singleton ground-state projector!,

$$\begin{aligned} E \star e^{-4E} &= e^{-4E} \star E = \frac{1}{2}e^{-4E}, \\ L_r^- \star e^{-4E} &= 0 = e^{-4E} \star L_r^+, \\ M_{rs} \star e^{-4E} &= 0 \end{aligned} \quad \Rightarrow \quad 4e^{-4E} \simeq |1/2; 0\rangle\langle 1/2; 0| \in \mathcal{D}_0 \otimes \mathcal{D}_0^*$$

(C.I., P. Sundell '08)

- General **spherically symm.** type-D sol.ns include *all* projectors on scalar (super)singleton modes (all $\mathfrak{so}(3)$ -invariant excitations of 4 $\exp(-4E)$) and their negative-energy counterparts.

$$\mathcal{P}_n(E) \sim \begin{cases} a^{\dagger i_1} \dots a^{\dagger i_n} \star |1/2; 0\rangle\langle 1/2; 0| \star a_{i_1} \dots a_{i_n}, & n > 0 \\ a_{i_1} \dots a_{i_{|n|}} \star |-1/2; 0\rangle\langle -1/2; 0| \star a^{\dagger i_1} \dots a^{\dagger i_{|n|}}, & n < 0 \end{cases}$$

$$a_1 = \frac{1}{2}(y_1 + iy_2), \quad a^{\dagger 1} = \frac{1}{2}(\bar{y}_1 - iy_2),$$

$$a_2 = \frac{1}{2}(-y_2 + iy_1), \quad a^{\dagger 2} = \frac{1}{2}(-\bar{y}_2 - iy_1)$$

$$[a_i, a^{\dagger j}]_{\star} = \delta_i^j$$

Spherically symmetric type-D solutions

- Using the gauge function:

$$\mathcal{P}_1(Y) = 4e^{-\frac{1}{2}Y^\alpha K'_{\alpha\beta} Y^\beta} = 4e^{-y^\alpha \sigma_{\alpha\dot{\alpha}}^0 \bar{y}^{\dot{\alpha}}} = 4e^{-4E} \rightarrow L^{-1} \star \mathcal{P}_1(Y) \star L = 4e^{-\frac{1}{2}Y^\alpha K_{\alpha\beta}(x) Y^\beta}$$

In AdS₄ spherical coords. (t,r,θ,φ) [ds² = (1+r²) dt² + (1+r²)⁻¹ dr² + r² dΩ²]

$$K'_{\alpha\beta} = (\Gamma_0)_{\alpha\beta} = \begin{pmatrix} 0 & u_\alpha^+ \bar{u}_\beta^+ + u_\alpha^- \bar{u}_\beta^- \\ \bar{u}_\alpha^+ u_\beta^+ + \bar{u}_\alpha^- u_\beta^- & 0 \end{pmatrix} \rightarrow K_{\alpha\beta} = \begin{pmatrix} 2r \tilde{u}_{(\alpha}^+ \tilde{u}_{\beta)}^- & \sqrt{1+r^2} (\tilde{u}_\alpha^+ \tilde{u}_\beta^+ + \tilde{u}_\alpha^- \tilde{u}_\beta^-) \\ \sqrt{1+r^2} (\tilde{u}_\alpha^+ \tilde{u}_\beta^+ + \tilde{u}_\alpha^- \tilde{u}_\beta^-) & 2r \tilde{u}_{(\alpha}^+ \tilde{u}_{\beta)}^- \end{pmatrix}$$

$$u^{+\alpha} u_\alpha^- = 1 = \tilde{u}^{+\alpha}(x) \tilde{u}_\alpha^-(x), \quad \varkappa^2(x) = -r^2$$

$$\Phi(x|Y) = \sum_n \nu_n \mathcal{N}_n \oint_{C(\varepsilon)} \frac{d\eta}{2\pi i} \left(\frac{\eta+1}{\eta-1} \right)^n \underbrace{L^{-1}(x) \star e^{-4\eta E} \star L(x) \star \kappa_y}$$

$$\bar{y} = 0 : \frac{1}{\eta r} \exp\left(\frac{1}{2\eta r} y^\alpha \tilde{u}_\alpha^+ \tilde{u}_\beta^- y^\beta\right) \Rightarrow \Phi_{\alpha(2s)}^{(n)} \sim \frac{i^{n-1} \mu_n}{r^{s+1}} (\tilde{u}^+ \tilde{u}^-)_{\alpha(2s)}^s$$

- Deformation parameter is real for scalar singleton, imaginary for spinor singl.
→ generalized electric/magnetic charge (or mass/NUT).

E/m duality connects Type A/B models?

- Spacetime coords. enter as parameter of a limit representation of a delta function . $\hat{\Phi}_1 \xrightarrow{r \rightarrow 0} \hat{\Phi}'_1 = \nu_1 \kappa_{y-i\sigma_0 \bar{y}} = 2\pi \nu_1 [\delta^2(y - i\sigma_0 \bar{y})]$

Cylindrically-symmetric type-D solutions

- Condition $K'_{\underline{\alpha}}{}^{\underline{\beta}} K'_{\underline{\beta}}{}^{\underline{\gamma}} = -\delta_{\underline{\alpha}}{}^{\underline{\gamma}}$ solved by any $Y^{\underline{\alpha}} K'_{\underline{\alpha}\underline{\beta}} Y^{\underline{\beta}} \sim E, J, iB, iP$

→ Solutions with $\mathfrak{so}(2,1)_{\mathfrak{h}} \oplus \mathfrak{so}(2)_{YK'Y}$ symmetry (centralizer of $YK'Y$).

- In particular, for $K' = \Gamma_{12}$, $\mathcal{P}_1(Y) := 4e^{-\frac{1}{2}Y^{\underline{\alpha}} K'_{\underline{\alpha}\underline{\beta}} Y^{\underline{\beta}}} = 4e^{-4J}$

Again a ground state of a 2D Fock-space (a non-compact ultra-short irrep, singleton-like but with roles of E and J exchanged, $|E| < |J|$ instead of $|E| > |J|$).
[Systematic procedure to extract creation/annihilation operators]

- Same steps yield $\Phi(x|Y) = \sum_n \nu_n \mathcal{N}_n \oint_{C(\varepsilon)} \frac{d\eta}{2\pi i} \left(\frac{\eta+1}{\eta-1} \right)^n \underbrace{L^{-1}(x) \star e^{-4\eta J} \star L(x) \star \kappa_y}$

$$\bar{y} = 0 : \frac{1}{\eta\sqrt{\varkappa^2}} \exp\left(\frac{1}{2\eta} y^{\alpha} \varkappa_{\alpha\beta}^{-1} y^{\beta}\right), \quad \varkappa_{\alpha\beta}^{-1} = -\frac{\varkappa_{\alpha\beta}}{\varkappa^2}, \quad \varkappa^2 = 1 + r^2 \sin^2 \theta$$

$$\Phi_{\alpha(2s)}^{(n)} \sim \frac{i^{n+s+1} \mu_n}{(1 + r^2 \sin^2 \theta)^{(s+1)/2}} (\tilde{u}^+ \tilde{u}^-)_{\alpha(2s)}^s$$

HS Invariants

- Define HS observables, gauge invariant off-shell. Weyl-curvature invariants:

$$\mathcal{C}_{2p}^{\pm} = \mathcal{N}_{\pm} \widehat{\text{Tr}}_{\pm}[\mathcal{C}_{2p}], \quad \mathcal{C}_{2p} = [\widehat{\Phi} \star \pi(\widehat{\Phi})]^{*p}$$

$$\widehat{\text{Tr}}_{+}[f(Y, Z)] = \int \frac{d^4 Y d^4 Z}{(2\pi)^4} f(Y, Z), \quad \widehat{\text{Tr}}_{-}[f(Y, Z)] = \widehat{\text{Tr}}_{+}[f(Y, Z) \star \kappa \bar{\kappa}]$$

- Ciclicity: $\widehat{\text{Tr}}_{\pm}[f(Y, Z) \star g(Y, Z)] = \widehat{\text{Tr}}_{\pm}[g(\pm Y, \pm Z) \star f(Y, Z)]$

- Conserved on the field equations:

$$\widehat{D}_{\mu} \widehat{\Phi} = 0 \Rightarrow \partial_{\mu} (\widehat{\Phi} \star \kappa)^{*q} = -[\widehat{A}_{\mu}, (\widehat{\Phi} \star \kappa)^{*q}]_{\star}$$

Ciclicity + A_{μ} even function of oscillators



$$d \widehat{\text{Tr}}_{\pm}[\mathcal{C}_{2p}^{\pm}] = 0$$

$$\mathcal{C}_k^{[0]} = \widehat{\text{Tr}}_{+} \left[(\widehat{\Phi} \star \pi(\widehat{\Phi}))^{*k} \star \kappa \bar{\kappa} \right]$$

$$\mathcal{I}(\sigma, k, \bar{k}; \lambda, \bar{\lambda}) = \widehat{\text{Tr}}_{+} \left[(\widehat{\kappa} \widehat{\bar{\kappa}})^{* \sigma} \star \exp_{\star}(\lambda^{\alpha} \widehat{S}_{\alpha} + \bar{\lambda}^{\dot{\alpha}} \widehat{S}_{\dot{\alpha}}) \star (\widehat{\Phi} \star \widehat{\kappa})^{*k} \star (\widehat{\Phi} \star \widehat{\bar{\kappa}})^{* \bar{k}} \right]$$

Singularity?

- Radial dependence of individual spin-s Weyl tensor $\sim r^{-s-1}$. However, HS-invariants for finitely many projectors are finite!

$$Tr_+ \left[(\widehat{\Phi} \star \pi(\widehat{\Phi}))^N \star \kappa \bar{\kappa} \right] = -4 \sum_{n=\pm 1, \pm 2, \dots} |n| (-1)^{(N+1)n} \mu_n^{2N}$$

Note: invariants are also (formally) insensitive to changes of ordering!
 Can the singularity be only an artefact of basis choice for function of operators?
 (crucial with non-polynomial f(operators))

- Examine master-fields in $r = 0$:

$$\Phi(r = 0) = L^{-1} |_{r=0} \star P_1(E) \star L |_{r=0} \star \kappa_y = P_1(E) \star \kappa_y \sim \delta^2(y - i\sigma^0 \bar{y})$$

$$\downarrow$$

$$[L(r = 0) = f(E)]$$

\Rightarrow Weyl tensors generating function $\sim \delta^2(y)$
 \rightarrow a regular function ($\exp(-2N_y)$) in normal ordering!

Conclusions & Outlook

- Found a general class of (almost) type-D solutions, with various symmetries:
 - spherical, HS generalization of Schwarzschild bh
 - cylindrical, HS counterpart of GR Melvin solution (regular everywhere)
 - biaxial (building blocks of the previous two, “almost type-D”)
 and other ones whose physical interpretation and GR analogues are yet to be studied.

- Singular? Not obvious, not at the level of invariants nor master-fields.
 - 1) A closed 2-form charge could detect singularities
 - 2) Divergent curvature invariants with infinitely many excitations
 A HS-invariant characterization of bhs is yet to be found.

- Must gain a better understanding of HS invariants and evaluate more of them.

[HS “metrics” $G_{\mu_1 \dots \mu_s} = \widehat{T}r_+ \left[\widehat{\kappa} \widehat{\kappa} \star \widehat{E}_{(\mu_1} \star \dots \star \widehat{E}_{\mu_s)} \right]$, $\widehat{E}_\mu = \frac{1}{2}(1 - \pi) \widehat{W}_\mu$]

- Multi-body solutions? [Preliminary analysis of consistency of a 2-body problem by evaluating 0-form invariants for $\Phi(x) + \Phi(y)$. Cross terms fall off as $V((1+r^2)^{-1/2}; n)$. Hierarchy of excitations ?]

- Thermodynamics in invariants? Horizon? Trapped surfaces?...

Conclusions & Outlook

- Study the boundary duals of such solutions. Many interesting questions:
 - What are the dual configurations in $U(N)/O(N)$ vector models?
 - Hawking-Page phase transitions? (*Shenker-Yin '11* \rightarrow *No uncharged bhs in Type A minimal model*)
 - Are spacetime boundary conditions (partly) encoded in (Y,Z) -space behaviour? [Distinction small/large gauge transformation and superselection sectors]
- Role of Z -space in non-perturbative sector of the theory . In particular, “ Z -space vacua”, topologically non-trivial flat Z -connections.
- Solutions mixing AdS massless particle state + soliton-like state. [Particles alone are inconsistent as solutions of the full eqs., backreaction forces addition of non-perturbative states]

Internal Z-Space Solution

- Ansatz for internal eqs., **separation of Y and Z** variables, absorb Y-dep. in $P_n(Y)$:

$$\widehat{S}'_\alpha = z_\alpha - 2i \sum_{n=0}^{\infty} P_n(Y) \star A_\alpha^n(z), \quad \widehat{\bar{S}}'_\alpha = \bar{z}_\alpha - 2i \sum_{n=0}^{\infty} P_n(Y) \star \bar{A}_\alpha^n(\bar{z})$$

Reduced deformed oscillators: $\Sigma_\alpha^n = z_\alpha - 2iA_\alpha^n, \quad \bar{\Sigma}_\alpha^n = \bar{z}_\alpha - 2i\bar{A}_\alpha^n$

- ✓ Orthogonality of projectors \Rightarrow eqs. for different n split;
- ✓ Projectors only Y-dep. \Rightarrow spectators, out of commutators;
- ✓ $\nu_n = \text{const}$ and $\pi(\Sigma) = -\Sigma$ solve $\{S', \Phi'\}_\pi = 0$;
- ✓ Holomorphicity in z of S' solves $[S', \bar{S}'] = 0$

- Left with the **deformed oscillator problem** :

$$\begin{aligned} [\Sigma_\alpha^n, \Sigma_\beta^n]_\star &= -2i\epsilon_{\alpha\beta}(1 - \mathcal{B}_n\nu_n\kappa_z), \\ [\bar{\Sigma}_\alpha^n, \bar{\Sigma}_\beta^n]_\star &= -2i\epsilon_{\alpha\beta}(1 - \bar{\mathcal{B}}_n\bar{\nu}_n\bar{\kappa}_{\bar{z}}) \end{aligned}$$

Can solve by a general method (*Prokushkin-Vasiliev '98, Sezgin-Sundell '05*) for regular deformation terms. Use a limit representation of $\kappa_z \sim \delta^2(z)$ or first go to normal-ordering where $\kappa_z = \text{gaussian}$.

Solution for Z-space deformed oscillators

- Introduce basis spinors u^\pm_α (a priori non-collinear with $\tilde{u}^\pm_\alpha(x)$) :

$$z^\pm := u^{\pm\alpha} z_\alpha, \quad w_z := z^+ z^-, \quad [z^-, z^+]_\star = -2i$$

- Solve $[\Sigma_n^+, \Sigma_n^-]_\star = -1 + \mathcal{B}_n \nu_n \kappa_z$ w/ Laplace-like transform:

$$\Sigma_n^\pm = 4z^\pm \int_{-1}^1 \frac{dt}{(t+1)^2} f_{\sigma_n}^{n\pm}(t) e^{i\sigma_n \frac{t-1}{t+1} w_z}$$

and using the limit representation $\lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} e^{-i\frac{\sigma}{\varepsilon} z^+ z^-} = \sigma [\kappa_z]^{\text{Weyl}}$.

Leads to manageable algebraic eqns for $f_{\pm}^n(t)$. Can either solve symmetrically, $f_+^n = f_-^n$, or asymmetrically (gauge freedom on S).

Study sym case: particular, ν -dependent solution

$$f_{\sigma_n}^{n\pm}(t) = \delta(t-1) - \frac{\sigma_n \mathcal{B}_n \nu_n}{4} {}_1F_1 \left[\frac{1}{2}; 2; \frac{\sigma_n \mathcal{B}_n \nu_n}{2} \log \frac{1}{t^2} \right]$$

- Also: a general way of solving the homogeneous ($\nu_n = 0$) eq. is the

projector solution: $X^2 = 1 \rightarrow X = 1 - 2P, \quad P^2 = P$

Solution for Z-space deformed oscillators

- Internal Z-space connection:

$$\begin{aligned}
 A_{\pm}^n &= A_{\pm}^{n(reg)} + A_{\pm}^{n(proj)} \\
 A_{\pm}^{n(reg)} &= \frac{i\sigma_n \mathcal{B}_n \nu_n}{2} z^{\pm} \int_{-1}^1 \frac{dt}{(t+1)^2} e^{i\sigma_n \frac{t-1}{t+1} w_z} \left[{}_1F_1 \left(1/2; 2; \frac{\nu_n}{2} \log \frac{1}{t^2} \right) \right] \\
 A_{\pm}^{n(proj)} &= -iz^{\pm} \sum_{k=0}^{\infty} (-1)^k \theta_k L_k[\nu_n] P_k(z), \quad P_k(z) = \frac{(z^+ z^-)^k}{k!} e^{-z^+ z^-}, \\
 L_k[\nu] &= \int_{-1}^1 dt t^k f_{\pm}^n(t) \longrightarrow 1 \text{ as } \nu_n \longrightarrow 0, \quad \theta_k = 0, 1
 \end{aligned}$$

Sol.ns depend on two infinite sets of parameters:

- continuous parameters $\nu_n \rightarrow \Phi$ -moduli;
 - discrete parameters $\theta_k \rightarrow S$ -moduli, a “landscape” of vacua.
- Divergent deformed oscillators ($t = -1$) but $S(x|Y,Z)$ only singular in $r = 0$!
 Pushed out of integration domain by star-product with $\mathcal{P}_n(x|Y)$. For $n=1$:

$$\widehat{S}^{\pm} = \tilde{z}^{\pm} + 8 \mathcal{P}_1(x|Y) \tilde{a}^{\pm} \int_{-1}^1 \frac{dt}{(t+1+i\sigma_n r(t-1))^2} j_1^{\pm}(t) e^{\frac{i\sigma_n(t-1)}{t+1+i\sigma_n r(t-1)}} \tilde{a}^+ \tilde{a}^-$$

$$\tilde{a}^{\pm} := \tilde{u}^{\alpha\pm} a_{\alpha}, \quad a_{\alpha} = z_{\alpha} + i(\kappa_{\alpha}^{\beta} y_{\beta} + \nu_{\alpha}^{\beta} \bar{y}_{\beta}), \quad z_{\alpha} \star \mathcal{P}_1 = a_{\alpha} \mathcal{P}_1, \quad [a_{\alpha}, a_{\beta}]_{\star} = -2i\epsilon_{\alpha\beta}$$

Exact solutions: gauge function method

- Y x Z-space eqns:

$$\begin{aligned}
 \widehat{F}_{\mu\nu} &= \widehat{F}_{\mu\alpha} = \widehat{F}_{\mu\dot{\alpha}} = 0, & \widehat{D}_\mu \widehat{\Phi} &= 0, \\
 [\widehat{S}'_\alpha, \widehat{S}'_\beta]_\star &= -2i\epsilon_{\alpha\beta}(1 - \mathcal{B} \star \widehat{\Phi}' \star \kappa), \\
 [\widehat{S}'_{\dot{\alpha}}, \widehat{S}'_{\dot{\beta}}]_\star &= -2i\epsilon_{\dot{\alpha}\dot{\beta}}(1 - \bar{\mathcal{B}} \star \widehat{\Phi}' \star \bar{\kappa}) \\
 [\widehat{S}'_\alpha, \widehat{S}'_{\dot{\beta}}]_\star &= 0, \\
 \widehat{S}'_\alpha \star \widehat{\Phi}' + \widehat{\Phi}' \star \pi(\widehat{S}'_\alpha) &= 0, \\
 \widehat{S}'_{\dot{\alpha}} \star \widehat{\Phi}' + \widehat{\Phi}' \star \bar{\pi}(\widehat{S}'_{\dot{\alpha}}) &= 0
 \end{aligned}$$

- Project on \mathcal{Z} ! (base \leftrightarrow fiber evolution)

Locally give x-dep. via gauge functions (spacetime \sim pure gauge!)

$$\widehat{A}_\mu = \widehat{L}^{-1} \star \partial_\mu \widehat{L}, \quad \widehat{S}_\alpha = \widehat{L}^{-1} \star (\widehat{S}'_\alpha) \star \widehat{L}, \quad \widehat{\Phi} = \widehat{L}^{-1} \star \widehat{\Phi}' \star \pi(\widehat{L})$$

$$\widehat{L} = \widehat{L}(x|Z, Y), \quad \widehat{L}(0|Z, Y) = 1 \quad \widehat{S}'_\alpha = \widehat{S}_\alpha(0|Z, Y), \quad \widehat{\Phi}' = \widehat{\Phi}(0|Z, Y)$$

- Z-eq.^{ns} can be solved exactly: 1) imposing **symmetries** on primed fields
2) via **projectors**
- “Dress” with x-dependence. Lorentz tensors are coefficients of:

$$\widehat{W}_\mu := \widehat{A}_\mu - \widehat{K}_\mu, \quad \widehat{K}_\mu := \frac{1}{4i} \omega_\mu^{\alpha\beta} \widehat{M}_{\alpha\beta} - \text{h.c.}, \quad \widehat{M}_{\alpha\beta} := y_\alpha y_\beta - z_\alpha z_\beta + \widehat{S}_{(\alpha} \star \widehat{S}_{\beta)}$$

Deformation parameters and asymptotic charges

- Building solutions on more than one projector opens up interesting possibilities.
- Every singleton-state projector contains a tower of fields of all spins \rightarrow can change basis and diagonalize on spin (and not occupation number)

$$\mathcal{C}(x|y) = \sum_n \nu_n \mathcal{N}_n \oint_{C(\epsilon)} \frac{d\eta}{2\pi i} \left(\frac{\eta+1}{\eta-1} \right)^n \frac{1}{\eta \sqrt{\kappa^2}} \exp \left(\frac{1}{2\eta} y^\alpha \kappa_{\alpha\beta}^{-1} y^\beta \right)$$

$$\mathcal{C}(x|y) = \sum_{s=0}^{\infty} \frac{1}{(2s)!} C_{\alpha(2s)}(x) y^{\alpha(2s)}, \quad C_{\alpha(2s)} \sim \frac{\mathcal{M}_s}{r^{s+1}} (\tilde{u}^+ \tilde{u}^-)_{\alpha(2s)}$$

“HS asymptotic charge”, $f(\nu_n)$:

$$\mathcal{M}_s \sim \sum_n \nu_n \tilde{\mathcal{N}}_n \oint_{C(\epsilon)} \frac{d\eta}{2\pi i \eta^{s+1}} \left(\frac{\eta+1}{\eta-1} \right)^n$$

(Can we choose ν_n such that $\mathcal{M}_s \sim \delta_{s,k}$, switching off all spins except one?)

- Possible to turn on an angular dependence in the Weyl tensor singularity via specific choices of deformation parameters (e.g. $\nu_n = q^n$, exchanging sum and integral) \rightarrow Kerr-like HS black-hole?

Reading off asymptotic charges

- Having found the gauge-fields generating functions, one may try to read off asymptotic charges from the sources of field strengths for $r \rightarrow \infty$, i.e. analyzing the asymptotics of the gauge field eq.

$$\nabla \widehat{W} + \widehat{W} \star \widehat{W} + \frac{1}{4i} \left(r^{\alpha\beta} \widehat{M}_{\alpha\beta} + \bar{r}^{\dot{\alpha}\dot{\beta}} \widehat{M}_{\dot{\alpha}\dot{\beta}} \right) = 0$$

$$r^{\alpha\beta} := d\omega^{\alpha\beta} + \omega^{\alpha\gamma} \omega^{\beta}_{\gamma}, \quad \nabla \widehat{W} = d\widehat{W} + \frac{1}{4i} \left[\omega^{\alpha\beta} \widehat{M}_{\alpha\beta}^{(0)} + \bar{\omega}^{\dot{\alpha}\dot{\beta}} \widehat{M}_{\dot{\alpha}\dot{\beta}}^{(0)}, \widehat{W} \right]_{\star}$$

after moving to the standard gauge of perturbation theory and reducing to spacetime submanifold $\{Z=0\}$.

- Possible mixing between different orders in \mathcal{M}_s due to s-dependent r-behaviour of spin-s component fields

$$(\nabla_{(0)} W + \{e_{(0)}, W\}_{\star})_{\alpha(2s)} \sim e_{(0)} \wedge e_{(0)} \partial_{\alpha(2s)}^{(y)} \Phi + \text{h.o.t.} = \frac{\widehat{\mathcal{M}}_s}{r^{s+1}} (u^+ u^-)_{\alpha(2s)}^s$$

leads to possible asymptotic charge redefinition

$$\widehat{\mathcal{M}}_s = \mathcal{M}_s + O(\mathcal{M}_s^2)$$

Twistor gauge and asymptotic charges

- To compare solutions in x-space, need to bring them in “universal” twistor gauge via some extended HS gauge transformation $G_{(v)}^{(K)}(x|Y,Z)$.

$$\widehat{v}^\alpha(x|Y, Z) \widehat{A}_\alpha(x|Y, Z) = \widehat{f}(x|Y, Z), \quad \frac{\partial}{\partial \nu_n} \widehat{v}^\alpha = \frac{\partial}{\partial K_{\alpha\beta}} \widehat{v}^\alpha = 0 = \frac{\partial}{\partial \nu_n} \widehat{f} = \frac{\partial}{\partial K_{\alpha\beta}} \widehat{f}$$

with residual gauge symmetries $\rightarrow \mathfrak{ho}(3,2)$, e.g., standard choice $v^\alpha = z^\alpha$.

- Our solutions are in some twistor gauge but NOT in universal twistor gauge (v^α depends explicitly on K). Can be brought to twistor gauge, e.g., the standard gauge of perturbative analysis, order by order in ν_n .
- The action of G_K^v on solutions will redefine the HS asymptotic charges, too!

$$\begin{aligned} \widehat{\Phi}_{(v)} &= (\widehat{G}_{(v)}^{(K)})^{-1} \star \widehat{\Phi}_{(K)} \star \pi(\widehat{G}_{(v)}^{(K)}) \\ \longrightarrow \mathcal{M}_s |_{(v)} &= \mathcal{M}_s |_{(K)} + \sum_{s, s''} \mathcal{M}_{s'} |_{(K)} \mathcal{M}_{s''} |_{(K)} f_s^{s' s''} + \dots \end{aligned}$$

- Finally, $\mathfrak{ho}(3,2)$ asymptotic symmetries (possibly enhanced to current algebra of free fields) will act \mathcal{M}_s . Invariants $\mathcal{O}(\mathcal{M}_s)$?